

DISCLAIMER: While Genzyme allows its Product Monograph to be downloaded and/or printed from its web site for your convenience, once this Product Monograph is downloaded and/or printed from its web site, Genzyme is unable to ensure the accuracy of the Product Monograph in the downloaded or printed version. The Product Monograph may be revised, updated and amended from time to time, and you are directed to the Product Monograph on the website (www.genzyme.ca) for the most current version.

PRODUCT MONOGRAPH

Pr **Aldurazyme[®]**

laronidase

**Solution for Intravenous Infusion
0.58 mg/mL (100 Units/mL)**

**Enzyme Replacement Therapy
ATC code: A16AB05**

Genzyme Canada Inc.
800-2700 Matheson Blvd. East, West Tower
Mississauga, ON
L4W 4V9
www.genzyme.ca

Date of Approval:
February 12, 2010

Submission Control No: 135931
Aldurazyme[®] is a registered trademark of BioMarin/Genzyme LLC

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	4
WARNINGS AND PRECAUTIONS	4
ADVERSE REACTIONS.....	6
DRUG INTERACTIONS	11
DOSAGE AND ADMINISTRATION	11
OVERDOSAGE	14
ACTION AND CLINICAL PHARMACOLOGY	14
PART II: SCIENTIFIC INFORMATION	17
PHARMACEUTICAL INFORMATION.....	17
CLINICAL TRIALS.....	19
DETAILED PHARMACOLOGY	21
TOXICOLOGY	23
REFERENCES	25
PART III: CONSUMER INFORMATION.....	27

Pr Aldurazyme®

laronidase

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form/ Strength	Clinically Relevant Nonmedicinal Ingredients
Intravenous infusion	Sterile solution/ 0.58 mg/mL (100 Units/mL) ¹	There are no clinically relevant nonmedicinal ingredients. Nonmedicinal ingredients: Sodium chloride, Sodium phosphate monobasic monohydrate, Sodium phosphate dibasic heptahydrate, Polysorbate 80, Water for Injection

¹Aldurazyme® is supplied as a sterile solution in clear Type I glass 5 mL vials (2.9 mg laronidase per 5 mL). The closure consists of a siliconized butyl stopper and an aluminum seal with a plastic flip-off cap.

INDICATIONS AND CLINICAL USE

Aldurazyme® (laronidase) is indicated for:

- long-term enzyme replacement therapy in patients with Mucopolysaccharidosis I (MPS I; α -L-iduronidase deficiency) to treat the non-central nervous system manifestations of the disease.

Pediatrics (6 months to 18 years of age): Aldurazyme® is indicated for enzyme replacement therapy in patients with Mucopolysaccharidosis I (MPS I; α -L-iduronidase deficiency) to treat the non-central nervous system manifestations of the disease.

Geriatrics (> 65 years of age): Clinical studies of Aldurazyme® did not include patients aged 65 years and over to determine whether they respond differently from younger patients.

CONTRAINDICATIONS

Patients who are severely hypersensitive to Aldurazyme[®] (laronidase) or to any ingredient in the formulation or component of the container (see **WARNINGS AND PRECAUTIONS: General**). For a complete listing of the ingredients in the formulation and components of the container, refer to the table in the **SUMMARY PRODUCT INFORMATION** section of the product monograph.

WARNINGS AND PRECAUTIONS

LIFE-THREATENING ANAPHYLACTIC REACTIONS HAVE BEEN OBSERVED IN SOME PATIENTS DURING ALDURAZYME[®] (LARONIDASE) INFUSIONS. THEREFORE, APPROPRIATE MEDICAL SUPPORT SHOULD BE READILY AVAILABLE WHEN ALDURAZYME IS ADMINISTERED. PATIENTS WITH COMPROMISED RESPIRATORY FUNCTION OR ACUTE RESPIRATORY DISEASE MAY BE AT RISK OF SERIOUS ACUTE EXACERBATION OF THEIR RESPIRATORY COMPROMISE DUE TO INFUSION REACTIONS, AND REQUIRE ADDITIONAL MONITORING.

General

Life-threatening anaphylactic reactions have been observed in some patients during or up to 3 hours after Aldurazyme[®] infusions. Reactions have included: respiratory failure, respiratory distress, stridor, tachypnea, bronchospasm, airway obstruction, hypoxia, hypotension, bradycardia, and urticaria. Interventions have included; resuscitation, mechanical ventilatory support, emergency tracheotomy, hospitalization, and treatment with inhaled beta-adrenergic agonists, epinephrine, and intravenous corticosteroids.

In clinical trials and post-marketing safety experience with Aldurazyme[®], approximately 1% of patients experienced severe or serious allergic reactions. In patients with MPS I, pre-existing upper airway obstruction may have contributed to the severity of some reactions. Due to the potential for severe allergic reactions, appropriate medical support should be readily available when Aldurazyme[®] is administered. Because of the potential for recurrent reactions, some patients who experience initial severe reactions may require prolonged observation.

Patients with an acute illness at the time of Aldurazyme[®] infusion may be at greater risk for infusion-related reactions. Careful consideration should be given to the patient's clinical status prior to administration of Aldurazyme[®].

Patients should receive antipyretics and/or antihistamines prior to infusion (see **ADVERSE REACTIONS**). If an infusion reaction occurs, regardless of pretreatment, decreasing the infusion rate, temporarily stopping the infusion, and/or administering additional antipyretics and/or antihistamines may ameliorate the symptoms (see **ADVERSE REACTIONS**). If

anaphylactic or other severe allergic reactions occur, immediately discontinue the infusion of Aldurazyme[®] and initiate appropriate treatment. Caution should be exercised if epinephrine is being considered for use in patients with MPS I due to the increased prevalence of coronary artery disease in these patients.

The risks and benefits of re-administering Aldurazyme[®] following an anaphylactic or severe allergic reaction should be considered. Extreme care should be exercised, with appropriate resuscitation measures available, if the decision is made to re-administer the product.

Carcinogenesis and Mutagenesis

Studies to assess the carcinogenic and mutagenic potential of Aldurazyme[®] have not been conducted.

Hepatic

The safety and efficacy of Aldurazyme[®] in patients with hepatic insufficiency have not been evaluated and no dosage regimen can be recommended in these patients.

Immune

Patients treated with Aldurazyme[®] may develop infusion-related allergic reactions (see **General** heading above and **ADVERSE REACTIONS**).

Renal

The safety and efficacy of Aldurazyme[®] in patients with renal insufficiency have not been evaluated and no dosage regimen can be recommended in these patients.

Special Populations

All patients should be informed that a registry for MPS I patients has been established and should be encouraged to participate in order to better understand and continue to monitor and evaluate MPS I disease and treatments. The MPS I Registry allows for the collection of data to further understand the long-term effectiveness and safety profile of Aldurazyme[®] treatment in this patient population. This should apply to those special populations listed below treated with Aldurazyme[®] (see **PART III, CONSUMER INFORMATION - MPS I Registry**). Information regarding the registry program may be found at www.MPSIRegistry.com or by calling (800) 745-4447.

Pregnant Women: No experience. Reproduction studies have been performed in male and female rats at doses up to 6.2 times the human dose and have revealed no evidence of impaired fertility or harm to the fetus due to Aldurazyme[®]. However, there are no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, Aldurazyme[®] should be used during pregnancy only if clearly needed. (See **TOXICOLOGY - Reproductive Toxicity**)

Nursing Women: It is not known whether the drug is excreted in human milk. Because many

drugs are excreted in human milk, caution should be exercised when Aldurazyme[®] is administered to a nursing woman.

Pediatrics (6 months to 18 years of age): Pediatric patients from 6 months up to 18 years of age have been treated with Aldurazyme[®] in clinical studies (see **ADVERSE REACTIONS** and **CLINICAL TRIALS** section). Long-term safety and efficacy data are not available.

Geriatrics (> 65 years of age): Clinical studies of Aldurazyme[®] did not include patients aged 65 years and over to determine whether they respond differently from younger patients.

Monitoring and Laboratory Tests

Evaluation of bioactivity during the clinical studies included changes in urinary glycosaminoglycan (GAG) levels, which were shown to decrease in patients treated with Aldurazyme[®] compared to those treated with placebo.

As seen in the clinical studies, it is expected that patients will develop antibodies to Aldurazyme[®]. It is strongly recommended that patients be monitored periodically for IgG antibody formation.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

The most serious adverse reactions reported with Aldurazyme[®] (laronidase) during clinical trials and the post-marketing period were anaphylactic and allergic reactions (see **WARNINGS AND PRECAUTIONS: General**, and **ADVERSE REACTIONS: Infusion-Related Reactions** and **Immunogenicity**).

The most common adverse reactions associated with Aldurazyme[®] treatment in the clinical studies were upper respiratory tract infection, rash, and injection site reaction (placebo-controlled study ALID-003-99), and pyrexia and chills (open-label study ALID-014-02).

The most common adverse reactions requiring intervention were infusion-related reactions. Most infusion-related reactions requiring intervention were ameliorated with slowing of the infusion rate, temporarily stopping the infusion, and/or administering additional antipyretics and/or antihistamines.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse

events and for approximating rates.

The following safety information is based on 2 clinical studies: a placebo-controlled study (ALID-003-99) and an open-label study (ALID-014-02). The data described in Table 1 reflect exposure to 0.58 mg/kg of Aldurazyme[®] for 26 weeks in a placebo-controlled double-blind study (ALID-003-99) in 45 patients with MPS I (N=22 Aldurazyme[®], and N=23 placebo). All 45 patients continued into an open-label study of Aldurazyme[®] treatment for an additional 36 weeks. An additional 10 patients participated in an early open-label study (BIO7500-001) with continued infusions for up to 3 years. The population in the placebo-controlled study (ALID-003-99) was evenly distributed for gender (N=23 females and 22 males) and ranged in age from 6 to 43 years. Of the 45 patients in the placebo-controlled study, 1 was clinically assessed as having the Hurler form, 37 Hurler-Scheie, and 7 Scheie. The data described in Table 2 reflect exposure to Aldurazyme[®] in an open-label study (ALID-014-02) of 20 patients 5 years of age or younger. Long-term safety data are not available given the 52-week duration of the study. The patients received weekly infusions of Aldurazyme[®] at a dose of 0.58 mg/kg for a total duration of 52 weeks, except for 4 patients who had their dose increased to 1.16 mg/kg from Week 26 onwards. Of the 20 patients, 16 were clinically assessed as having the Hurler form and 4 Hurler-Scheie. All patients were treated with antipyretics and antihistamines prior to the infusions.

Table 1 enumerates adverse events and selected laboratory abnormalities that occurred during the placebo-controlled trial in at least 2 patients more in the Aldurazyme[®] group than was observed in the placebo group. Reported adverse events have been classified using standard WHOART terms. Observed adverse events in the early open-label study (BIO7500-001) and the open-label treatment period following the controlled study (ALID-006-01) were not different in nature or severity.

Table 1: Number and (%) of Patients with Adverse Events and Selected Laboratory Abnormalities in the Placebo-Controlled Study (ALID-003-99)

Adverse Event	Aldurazyme[®] (N=22)	Placebo N=23
Respiratory System		
Upper respiratory tract infection	7 (32)	4 (17)
Body as a Whole		
Chest pain	2 (9)	0
Nervous System		
Hyperreflexia	3 (14)	0
Paresthesia	3 (14)	1 (4)
Skin and Appendages		
Rash	8 (36)	5 (22)
Resistance Mechanism		
Abscess	2 (9)	0
Liver and Biliary System		
Bilirubinemia	2 (9)	0
Vascular		
Vein disorder	3 (14)	1 (4)
Urinary System		
Facial edema	2 (9)	0
Cardiovascular, General		
Hypotension	2 (9)	0
Dependent edema	2 (9)	0
Vision		
Corneal opacity	2 (9)	0
Application Site		
Injection site pain	2 (9)	0
Injection site reaction	4 (18)	2 (9)
Platelet, Bleeding and Clotting		
Thrombocytopenia	2 (9)	0

Table 2 presents a summary of the adverse reactions that occurred during the open-label trial in at least 5% (1 or more) of the patients. Reported adverse reactions have been classified using MedDRA terminology.

Table 2: Summary of Adverse Events Related to Treatment in ≥ 5% Patients in Open-Label Study (ALID-014-02)

System Organ Class Preferred Term	0.58 mg/kg N=16 n (%)	0.58-1.16 mg/kg* N=4 n (%)	Overall N=20 n (%)
Any Adverse Events	5 (31)	3 (75)	8 (40)
Cardiac disorders	1 (6)	1 (25)	2 (10)
Tachycardia	1 (6)	1 (25)	2 (10)
General disorders and administration site conditions	5 (31)	3 (75)	8 (40)
Pyrexia	4 (25)	3 (75)	7 (35)
Chills	3 (19)	1 (25)	4 (20)
Crepitations	0	1 (25)	1 (5)
Investigations	1 (6)	1 (25)	2 (10)
Blood pressure increased	1 (6)	1 (25)	2 (10)
Oxygen saturation decreased	1 (6)	1 (25)	2 (10)
Blood iron decreased	0	1 (25)	1 (5)
Heart rate increased	1 (6)	0	1 (5)
Nervous system disorders	0	1 (25)	1 (5)
Tremor	0	1 (25)	1 (5)
Respiratory, thoracic and mediastinal disorders	0	1 (25)	1 (5)
Respiratory distress	0	1 (25)	1 (5)
Wheezing	0	1 (25)	1 (5)
Skin and subcutaneous tissue disorders	0	1 (25)	1 (5)
Pruritus	0	1 (25)	1 (5)
Rash macular	0	1 (25)	1 (5)
Vascular disorders	0	2 (50)	2 (10)
Hypertension	0	1 (25)	1 (5)
Pallor	0	1 (25)	1 (5)

Note: Percentages are based on the total number of patients in the dose group.
Note: A patient experiencing more than 1 adverse event within a system organ class or preferred term is counted once within that system organ class or preferred term.
Note: The dose from Week 26 onwards was increased to 1.16 mg/kg, if the urinary GAG level at Week 22 was > 200 µg/mg creatinine for the group of patients enrolled after 1 January 2004.
*1.16 mg/kg as of Week 26

Infusion-Related Reactions

Infusion-related reactions were reported in 7 of 22 patients treated with Aldurazyme[®] in the placebo-controlled study (ALID-003-99). Infusion-related reactions were not significantly different between the Aldurazyme[®] treatment group and the placebo (infusions of diluent and all nonmedicinal components of Aldurazyme[®]) group. The most common infusion-related reactions included flushing, fever, headache and rash. Flushing occurred in 5 patients (23%) receiving Aldurazyme[®]; the other reactions were less frequent. All reactions were mild to moderate in severity. The frequency of infusion-related reactions decreased with continued use during the open-label extended use period. There was one case of anaphylaxis during the open-label extension period (ALID-006-01) (see **WARNINGS AND PRECAUTIONS: General**). Less common infusion-related reactions include cough, bronchospasm, dyspnea, urticaria, angioedema and pruritus.

In the open-label study (ALID-014-02), infusion-related reactions were reported in a total of 7 (35%) of patients. The most frequently reported infusion-related reactions were pyrexia (6 patients) and chills (4 patients). The majority of infusion-related reactions were mild in severity, and could be managed with a decrease of the infusion rate, a temporary interruption of the infusion, and/or administration of antihistamine and/or antipyretic.

Immunogenicity

The data reflect the percentage of patients whose test results were considered positive for antibodies to Aldurazyme[®] using an enzyme-linked immunosorbent assay (ELISA) for laronidase-specific IgG binding antibodies, and are highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibodies in an assay may be influenced by several factors including sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies to Aldurazyme[®] with the incidence of antibodies to other products may be misleading.

In the placebo-controlled study (ALID-003-99) and an early open-label study (BIO7500-001), 50 of 55 patients (91%) treated with Aldurazyme[®] were positive for IgG antibodies to laronidase. Similarly, 20 of 20 patients (100%) treated with Aldurazyme[®] in the open-label study (ALID-014-02) developed IgG antibodies to laronidase. One patient who initially developed IgG antibodies to laronidase no longer had detectable IgG antibodies after a total of 12 months of Aldurazyme[®] treatment in the open-label study (ALID-014-02). All seropositive patients in this open-label study (ALID-014-02) were tested for in-vitro neutralizing effects on enzyme activity. One patient showed marginal to low level in-vitro neutralizing inhibitory activity, which did not appear to impact clinical efficacy and/or urinary GAG reduction. The clinical significance of antibodies to Aldurazyme[®] is not known, including the potential for product neutralization.

Four patients in the placebo-controlled study (ALID-003-99) who experienced severe infusion-related reactions and 2 patients in the open-label study (ALID-014-02) who experienced moderate infusion related reactions were tested for Aldurazyme[®]-specific IgE antibodies and complement activation. IgE testing was performed by ELISA and complement activation was measured by the Quidel Enzyme Immunoassay. Only one of these patients (in the placebo-controlled study ALID-003-99) tested positive for both Aldurazyme[®]-specific IgE binding antibodies and complement activation; this patient had an anaphylactic reaction consisting of urticaria and airway obstruction (see **WARNINGS AND PRECAUTIONS: General**).

Other allergic reactions were also seen in patients receiving Aldurazyme[®] (see **ADVERSE REACTIONS: Infusion-Related Reactions**).

Post-Market Adverse Drug Reactions

In post-marketing experience with Aldurazyme[®], severe and serious infusion-related reactions have been reported, some of which were life-threatening (see **WARNINGS AND PRECAUTIONS, General**). The most common adverse reactions (using MedDRA terminology) included: chills, vomiting, nausea, arthralgia, diarrhea, tachycardia, abdominal pain, blood pressure increased, and oxygen saturation decreased. Additional adverse reactions identified in the post-marketing setting also include erythema, feeling cold and cyanosis.

There have been a small number of reports of extravasation in patients treated with ALDURAZYME. There have been no reports of tissue necrosis associated with extravasation.

DRUG INTERACTIONS

Drug-Drug Interactions: Interactions with other drugs have not been established. Physician monitoring is required for patients using medicinal products that contain chloroquine or procaine due to the potential risk of interference with the intracellular uptake of laronidase.

Drug-Food Interactions: Interactions with food have not been established.

Drug-Herb Interactions: Interactions with herbal products have not been established.

Drug-Laboratory Interactions: Interactions with laboratory tests have not been established.

DOSAGE AND ADMINISTRATION

Dosing Considerations

- Aldurazyme[®] (laronidase) treatment should be supervised by a physician experienced in the management of patients with MPS I or other inherited metabolic diseases.
- Administration of Aldurazyme[®] should be carried out in an appropriate clinical setting

- where resuscitation equipment to manage medical emergencies is readily available.
- The safety and efficacy of Aldurazyme[®] in patients older than 65 years have not been established and no dosage regimen can be recommended in these patients.
 - The safety and efficacy of Aldurazyme[®] in patients with renal or hepatic insufficiency have not been evaluated and no dosage regimen can be recommended in these patients.

Recommended Dose and Dosage Adjustment

The recommended dosage regimen of Aldurazyme[®] is 0.58 mg/kg body weight administered once weekly as an intravenous infusion.

Pretreatment with antipyretics and/or antihistamines is recommended 60 minutes prior to the start of the infusion (see **WARNINGS AND PRECAUTIONS: General**).

The initial infusion rate of 10 µg/kg/hr may be incrementally increased every 15 minutes during the first hour, as tolerated, until a maximum infusion rate of 200 µg/kg/hr is reached. The maximum rate is then maintained for the remainder of the infusion (2-3 hours).

The total volume of the infusion is determined by the patient's body weight and should be delivered over approximately 3 to 4 hours. Patients with a body weight of 20 kg or less should receive a total volume of 100 mL. Patients with a body weight of greater than 20 kg should receive a total volume of 250 mL.

The following tables provide infusion regimens and volumetric pump settings designed to administer the total volume over the recommended infusion time:

For Patients weighing 20 kg or Less

Total Volume of Aldurazyme[®] Infusion = 100 mL	
2 mL/hr x 15 minutes	Obtain vital signs, if stable then increase the rate to ...
4 mL/hr x 15 minutes	Obtain vital signs, if stable then increase the rate to ...
8 mL/hr x 15 minutes	Obtain vital signs, if stable then increase the rate to ...
16 mL/hr x 15 minutes	Obtain vital signs, if stable then increase the rate to ...
32 mL/hr x ~ 3 hours	For the remainder of the infusion

For Patients weighing Greater than 20 kg

Total Volume of Aldurazyme[®] Infusion = 250 mL	
5 mL/hr x 15 minutes	Obtain vital signs, if stable then increase the rate to ...
10 mL/hr x 15 minutes	Obtain vital signs, if stable then increase the rate to ...
20 mL/hr x 15 minutes	Obtain vital signs, if stable then increase the rate to ...
40 mL/hr x 15 minutes	Obtain vital signs, if stable then increase the rate to ...
80 mL/hr x ~ 3 hours	For the remainder of the infusion

Administration

Instructions for Use (With Aseptic Techniques and Sterile Preparation)

1. Determine the number of vials to be diluted based on the individual patient's weight and the recommended dose of 0.58 mg/kg:

$$\begin{aligned} \text{Patient's weight (kg)} \times 1 \text{ mL/kg of Aldurazyme}^{\text{®}} &= \text{Total \# mL of Aldurazyme}^{\text{®}} \\ \text{Total \# mL of Aldurazyme}^{\text{®}} \div 5 \text{ mL per Vial} &= \text{Total \# of Vials} \\ &\text{Round up to the nearest whole vial.} \end{aligned}$$

Remove the required number of vials from the refrigerator to allow them to reach room temperature. Do not heat or microwave the vials.

2. Before withdrawing the Aldurazyme[®] from the vial, visually inspect each vial for particulate matter and discoloration. The Aldurazyme[®] solution should be clear to slightly opalescent and colorless to pale yellow. A few translucent particles may be present. Do not use if the solution is discolored or if there is particulate matter in the solution.

3. Determine the total volume of the infusion to be used based on the patient's body weight. The total final volume should be either 100 mL (if patient weight is less than or equal to 20 kg) or 250 mL (if patient weight is greater than 20 kg).

4. Using the chart below, prepare an infusion bag of 0.1% Albumin (Human) in 0.9% Sodium Chloride Injection, USP. Remove and discard a volume of 0.9% Sodium Chloride Injection, USP equal to the volume of Albumin (Human) to be added to the infusion bag. Add the appropriate volume of Albumin (Human) to the infusion bag and gently rotate the infusion bag to ensure proper distribution of the Albumin.

Total Volume of Aldurazyme[®] Infusion	Volume of Albumin (Human) 5% to be Added	Volume of Albumin (Human) 25% to be Added
100 mL	2 mL	0.4 mL
250 mL	5 mL	1 mL

5. Withdraw and discard a volume of the 0.1% Albumin (Human) in 0.9% Sodium Chloride Injection, USP from the infusion bag, equal to the volume of Aldurazyme[®] concentrate to be added.

6. Slowly withdraw the calculated volume of Aldurazyme[®] from the appropriate number of vials using caution to avoid excessive agitation. Do not use a filter needle, as this may cause agitation. Agitation may denature Aldurazyme[®], rendering it biologically inactive.

7. Slowly add the Aldurazyme[®] solution to the 0.1% Albumin (Human) in 0.9% Sodium Chloride Injection, USP using care to avoid agitation of the solutions. Do not use a filter needle.

8. Gently rotate the infusion bag to ensure proper distribution of Aldurazyme[®]. Do not shake the solution.

Aldurazyme[®] does not contain any preservatives; therefore, after dilution with saline in the infusion bags, any unused product or waste material should be discarded and disposed of in accordance with local requirements.

Aldurazyme[®] must not be mixed with other medicinal products in the same infusion.

The compatibility of Aldurazyme[®] in solution with other products has not been evaluated.

Storage and Stability

Store Aldurazyme[®] (laronidase) under refrigeration at 2 to 8°C (36 to 46°F). DO NOT FREEZE OR SHAKE. DO NOT USE Aldurazyme[®] after the expiration date on the vial. This product contains no preservatives.

It is recommended that once Aldurazyme[®] is diluted, the infusion should start immediately (within 3 hours) as there is no preservative in either Aldurazyme[®] or in the infusion bag. Although not recommended, physicochemical stability studies have shown that the diluted solution may be stored at 2 to 8°C for up to 36 hours if aseptic technique is used throughout the procedure.

Special Handling Instructions

Do not use the vial more than one time. Aldurazyme[®] should be prepared using PVC containers and administered with a PVC infusion set equipped with an in-line, low-protein binding 0.2 micrometer (µm) filter. There is no information on the compatibility of diluted Aldurazyme[®] with glass containers.

OVERDOSAGE

There have been no reports of overdose with Aldurazyme[®].

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Mucopolysaccharide storage disorders are caused by the deficiency of specific lysosomal enzymes required for the catabolism of glycosaminoglycans (GAGs). Mucopolysaccharidosis I (MPS I) is characterized by the deficiency of α -L-iduronidase, a lysosomal hydrolase which catalyses the hydrolysis of terminal α -L-iduronic acid residues of dermatan sulfate and heparan sulfate. Reduced or absent α -L-iduronidase activity results in the accumulation of the GAG substrates, dermatan sulfate and heparan sulfate, throughout the body and leads to widespread cellular, tissue, and organ dysfunction.

The rationale of Aldurazyme[®] (laronidase) therapy in MPS I is to provide exogenous enzyme for uptake into lysosomes and increase the catabolism of GAG. Aldurazyme[®] uptake by cells into lysosomes is most likely mediated by the mannose-6-phosphate-terminated oligosaccharide chains of laronidase binding to specific mannose-6-phosphate receptors.

Pharmacodynamics

Because many proteins in the blood are restricted from entry into the central nervous system (CNS) by the blood brain barrier, effects of intravenously administered Aldurazyme[®] on cells within the central nervous system cannot be inferred from activity in sites outside the CNS. The ability of Aldurazyme[®] to cross the blood brain barrier has not been evaluated in animal models or in clinical trials.

Pharmacokinetics

The pharmacokinetics of laronidase were evaluated in 12 patients with MPS I who received 0.58 mg/kg of Aldurazyme[®] as a 4-hour infusion in the placebo-controlled study (ALID-003-99). After the 1st, 12th and 26th weekly infusions, the mean maximum plasma concentrations (C_{max}) ranged from 1.2 to 1.7 μ g/mL for the 3 time points. The mean area under the plasma concentration-time curve (AUC) ranged from 4.5 to 6.9 μ g • hr/mL. The mean volume of distribution (V_z) ranged from 0.24 to 0.6 L/kg. Mean plasma clearance (CL) ranged from 1.7 to 2.7 mL/min/kg, and the mean elimination half-life ($t_{1/2}$) ranged from 1.5 to 3.6 hr.

The pharmacokinetic profile in the open-label study of 20 patients aged 5 years or younger (ALID-014-02) was similar to that of older, less severely affected patients (**Table 3**).

Table 3: Summary of Pharmacokinetic Parameters for Aldurazyme[®] at Weeks 1, 13, 26, and 52 (ALID-014-02)

Parameter	Week 1	Week 13	Week 26	Week 52
C_{max}^1 (μ g/mL)	1.89 \pm 3.88	0.90 \pm 0.53	2.15 \pm 2.33	3.13 \pm 5.12
AUC _{∞} ¹ (h* μ g/mL)	3.35 \pm 3.43	2.43 \pm 1.43	5.34 \pm 3.94	5.45 \pm 5.63
$t_{1/2}$ (h)	1.55 \pm 0.52	0.55 \pm 0.28	0.95 \pm 0.88	1.15 \pm 0.87
CL(mL/min/kg)	5.73 \pm 3.90	5.33 \pm 2.67	2.97 \pm 1.87	3.20 \pm 2.02
V_z (L/kg)	0.75 \pm 0.50	0.30 \pm 0.30	0.21 \pm 0.30	0.25 \pm 0.21
V_{ss} (L/kg)	0.89 \pm 0.69	0.70 \pm 0.40	0.42 \pm 0.28	0.48 \pm 0.31
Data are reported as mean \pm standard deviation.				
¹ Assuming an average potency of 172.4 Units of activity/mg protein				

Effects of Antibodies

Most patients who received once-weekly infusions of Aldurazyme[®] in the placebo-controlled study (ALID-003-99) developed antibodies to laronidase by week 12. Between weeks 1 and 12, increases in plasma clearance of laronidase were observed in some patients that appeared to be proportional to the antibody titer. At week 26, plasma clearance of laronidase was comparable to that at week 1, in spite of the continued and, in some cases, increased titers of antibodies.

In the open-label study (ALID-014-02), all patients developed antibodies to laronidase by Week 12. Aldurazyme[®] exposure was similar across visits but variable across patients. Taking into account the small sample size and variability, the effect of antibody titers on plasma clearance, maximum plasma concentration and area under the plasma concentration-time curve in patients of 5 years of age or younger was comparable to that observed in the older, less severely affected patients in the placebo-controlled study (ALID-003-99). There was an apparent decrease in volume of distribution which resulted in a decrease in half-life with continued dosing. This appears to be related to an increase in antibody titers.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: laronidase

Chemical name: recombinant human α -L-iduronidase (rhIDU)

Molecular formula and molecular mass: $C_{3169}H_{4889}O_{884}N_{901}S_{12}$

Structural formula:



Physicochemical properties: Recombinant human α -L-iduronidase (rhIDU) is a 628 amino acid lysosomal hydrolase that cleaves iduronic acid residues from the nonreducing ends of heparan sulfate and dermatan sulfate. The enzyme is a single polypeptide chain of molecular mass 70.1 kilodaltons (from translated cDNA sequence). This is a soluble monomeric protein with an apparent molecular weight by SDS-PAGE analysis, of 83 kD. rhIDU contains six asparagine-linked glycosylation sites, two of which carry the bis mannose-6-phosphate oligomannose₇ oligosaccharide that binds the target cell surface receptor.

Specific Activity: 172 U/mg

Product Characteristics

Aldurazyme[®], (laronidase) for intravenous infusion, is supplied as a sterile, nonpyrogenic, colorless to pale yellow, clear to slightly opalescent solution that must be diluted prior to administration in 0.9% Sodium Chloride Injection, USP, containing 0.1% Albumin (Human). The solution in each vial contains a nominal laronidase concentration of 0.58 mg/mL and a pH of approximately 5.5. The extractable volume of 5.0 mL from each vial provides 2.9 mg laronidase, 43.9 mg sodium chloride, 63.5 mg sodium phosphate monobasic monohydrate, 10.7 mg sodium phosphate dibasic heptahydrate, and 0.05 mg polysorbate 80. Aldurazyme[®] does not contain preservatives; vials are for single use only.

Viral Inactivation

The viral safety of Aldurazyme[®] is confirmed by a combination of selection and qualification of vendors, raw material testing, cell bank characterization studies, validation of the viral removal and inactivation capacity of the rhIDU purification process, and routine in-process testing.

CLINICAL TRIALS

Study demographics and trial design

Table 4:

Study #	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age (Range)	Gender
ALID-003-99	Randomized, placebo-controlled, double blind	Aldurazyme [®] at 0.58 mg/kg or placebo, intravenous infusion, once weekly for 26 weeks	MPS I patients (n = 45)	15.5 years (6 to 43 years)	22 M/ 23 F
ALID-014-02	Open-Label ¹	Aldurazyme [®] at 0.58 mg/kg, intravenous infusion, once weekly for 52 weeks ²	MPS I patients (n = 20)	2.9 years (0.5 to 5.1 years)	12 M/ 8 F

¹ Patients were excluded from ALID-014-02 trial if: the patient was under consideration for or had undergone Hematopoietic Stem Cell Transplantation (HSCT); the patient had acute hydrocephalus at the time of enrollment; the patient had a clinically significant organic disease (with the exception of symptoms relating to MPS I) including: cardiovascular, hepatic, pulmonary, neurologic, or renal disease, other serious intercurrent illness, or extenuating circumstances that, in the opinion of the Investigator, would preclude participation in the trial or potentially decrease survival; the patient had received any investigational product within 30 days prior to trial enrollment; the patient has known severe hypersensitivity to Aldurazyme[®] or components of the delivery solution.

² In 4 patients enrolled after January 1, 2004, the dose was increased from Week 26 onwards (to Week 52) to 1.16 mg/kg for a urinary GAG level at Week 22 that was > 200 µg/mg creatinine

Study results

Aldurazyme[®] was studied in a randomized, placebo-controlled clinical trial (ALID-003-99) of 45 MPS I patients of whom 1 patient was clinically assessed as having the Hurler form, 37 Hurler-Scheie, and 7 Scheie. All patients had a baseline forced vital capacity (FVC) less than or equal to 77% of predicted. Patients received Aldurazyme[®] at 0.58 mg/kg or placebo once-weekly for 26 weeks. All patients were treated with antipyretics and antihistamines prior to each infusion.

The primary efficacy outcome assessments were FVC and distance walked in 6 minutes (6-minute walk test, 6MWT). After 26 weeks, patients treated with Aldurazyme[®] showed improvement in FVC and in 6MWT compared to placebo-treated patients (see **Table 5**).

Table 5: Primary Efficacy Outcomes

	Aldurazyme[®] N = 22	Placebo N = 23
Forced Vital Capacity (percent of predicted normal)		
Baseline (Mean ± s.d.)	48 ± 15	54 ± 16
Week 26 (Mean ± s.d.)	53 ± 18	54 ± 14
Change from baseline to week 26 (Mean ± s.d.)	5 ± 9	-1 ± 6
Change from baseline to week 26 (median)	3	0
Difference between groups (mean)	6	
Difference between groups Median (95% CI)	3 p=0.009*	
6-Minute Walk Distance (metres)		
Baseline (Mean ± s.d.)	319 ± 131	367 ± 114
Week (Mean ± s.d.)	339 ± 127	348 ± 129
Change from baseline to week 26 (Mean ± s.d.)	20 ± 69	-18 ± 67
Change from baseline to week 26 (median)	28	-11
Difference between groups (mean)	38	
Difference between groups Median (95% CI)	39 (-2, 79) p=0.07*	

*By Wilcoxon Rank Sum Test


Evaluations of bioactivity were changes in liver size and urinary GAG levels. Liver size and urinary GAG levels decreased in patients treated with Aldurazyme[®] compared to patients treated with placebo. No subject in the group receiving Aldurazyme[®] reached the normal range for urinary GAG levels during this 6-month study.

All 45 patients received open-label Aldurazyme[®] for 36 weeks following the double-blind period (ALID-006-01). Maintenance of mean FVC and an additional increase in mean 6MWT distance were observed compared to the start of the open-label period among patients who were initially randomized to and then continued to receive Aldurazyme[®]. Among patients who had been initially randomized to placebo, improvements from baseline in mean FVC and 6MWT distance were observed compared to the start of the open-label period.

An open-label, 1-year study (ALID-014-02) was conducted that mainly assessed the safety and pharmacokinetics of Aldurazyme[®] in 20 patients less than 5 years of age (16 patients were of the Hurler form and 4 of the Hurler-Scheie form). Other objectives were to evaluate the efficacy of Aldurazyme[®] by assessing changes in liver size, urinary GAG excretion, upper respiratory care requirements, sleep apnea, hearing, vision, growth velocity, electrocardiogram (ECG), echocardiogram, and the investigator's global assessment. Long term efficacy data are not available given the 52-week duration of the study. The patients received Aldurazyme[®] 0.58 mg/kg weekly infusions for a total duration of 52 weeks, except for 4 patients who had their dose increased to 1.16 mg/kg from Week 26 onwards. Eighteen patients completed the study and 2 patients died as result of cardiac failure and respiratory arrest, respectively, which were unrelated to treatment but related to complications of MPS I disease.

Urinary GAG levels decreased in all patients compared to baseline. The mean percentage reduction in urinary GAG level was 61.3%. The liver volumes of all patients were classified as abnormally high at baseline. At Week 52, the liver volumes of 9 of the 18 patients who completed the study were classified as normal and the remainder had shown a decrease in liver size. Echocardiography showed the mean left ventricular mass to have decreased (of 10 patients with mild left ventricular hypertrophy at baseline, 7 patients normalised at Week 52) and mean ejection fraction had decreased but remained within normal range. Some valvular changes had occurred. Younger patients with severe MPS I and patients with intermediate severity MPS I showed growth rates (height and weight) which approached normal and improvements in mental development and adaptive behavior, whilst older children with severe MPS I did not.

Figure 1: Urinary GAG Levels ($\mu\text{g}/\text{mg}$ creatinine) versus Age (ITT population)



DETAILED PHARMACOLOGY

Summary of Pharmacodynamic Studies

The pharmacodynamics of rhIDU were studied *in vitro* and in dogs and cats with MPS I. The *in vitro* studies explored the uptake of rhIDU by MPS I patient fibroblasts, its effect on GAG storage and its half-life in the cells.

Seven animal studies (six canine and one feline) were conducted to assess the potential pharmacodynamic effects of rhIDU in the treatment of MPS I. The first three studies focused on short-, medium- and long-term treatment of MPS I dogs with a low dose of 0.1 mg/kg was given

1–3 times per week or weekly. Subsequent studies explored short- and then long-term treatment of MPS I dogs with higher doses of 0.5 mg/kg/week and ultimately a dose of 2.0 mg/kg/week given as a bolus or continuous intravenous infusion. The one study in MPS I cats explored the efficacy of doses of 0.1 and 0.5 mg/kg/week. Finally, a study was conducted in normal dogs to determine whether the diluent used to prepare the rhIDU infusate affected the tolerance and immune response of the dogs to rhIDU.

Overall, the results from the six efficacy studies in MPS I dogs indicated that a dose of at least 0.5 mg/kg/week of rhIDU was necessary to obtain measurable levels of enzyme activity in all tissues examined. This resulted in decreased GAG storage in many tissues, particularly in liver, kidney and spleen. Increasing the dose to 2.0 mg/kg/week resulted in increased tissue enzyme levels for C.I. and for bolus. However, the bolus dose of 0.5 mg/kg/week was adequate to result in the highest achievable reduction of GAG in most tissues. Results from these two studies indicate that 0.5 mg/kg/week is an adequate dose for long-term therapy in MPS I dogs. The half-life of rhIDU in fibroblasts from MPS I patients *in vitro* was shown to be 5 days, suggesting that the once-weekly dosing regimen should be adequate to maintain tissue levels.

Thus, studies of the efficacy of rhIDU in lowering tissue GAG, together with the half-life of the enzyme *in vitro*, support the recommended human dose of 100 U/kg (0.58 mg/kg) given weekly.

Treatment of MPS I dogs or cats with rhIDU was well tolerated with no adverse clinical, clinical pathology or histomorphological findings with one exception. Anaphylactoid reactions were observed during infusion of rhIDU in several early dog studies, as well as in the study in cats. These reactions, which could be severe, were managed successfully by stopping the infusion, administering IV fluids and, if necessary, oxygen. The probable cause of the reactions was shown to be IgG-mediated complement activation and in most subsequent studies animals were pretreated with antihistamines prior to infusions.

The anaphylactoid reactions observed in the early preclinical studies were not observed with the clinical drug product in either monkeys or dogs, even without pretreatment with antihistamine drugs or in the absence of serum albumin in the infusate.

Summary of Pharmacokinetic Studies

In one study, a female MPS I dog received 0.1 mg/kg on three consecutive days. rhIDU exhibited a biphasic clearance pattern from the plasma with a presumed distribution phase ($t_{1/2\alpha} = 0.9$ minutes) and a slow clearance phase ($t_{1/2\beta} = 18.9$ minutes). The clearance of free enzyme from the plasma was associated with uptake by peripheral leukocytes; levels peaked four hours after the dose.

In a second study, two MPS I dogs received a weekly intravenous 9-hour bolus infusion of 2.0 mg/kg every week for 10 weeks. During week 2 of treatment, rhIDU was cleared biphasically

from the plasma of the two dogs, but it was cleared monophasically during week 10 of treatment. At week 2, the initial decline in concentration was very rapid with α $t_{1/2}$ = 0.61 and 0.94 minutes; the second, terminal phase was slower with β $t_{1/2}$ = 59.5 and 84.9 minutes. At week 10, the concentration declined monophasically with terminal half-lives of 66.2 and 23.8 minutes, respectively, in the two dogs. The volume of distribution, V_c , approximately 60 mL/kg, indicated that the enzyme was distributed mostly in the plasma, which has a volume of approximately 35 mL/kg. The clearance (Cl) of enzyme activity appeared to decrease considerably from week 2 to week 10, and there was a concomitant increase in the AUC. The reasons for these changes are not known.

TOXICOLOGY

Carcinogenesis

Studies to assess the carcinogenic potential of rhIDU have not been conducted. Carcinogenic potential would not be anticipated with rhIDU based on the structure of the drug substance (a recombinant human glycoprotein) and its impurity profile. The biochemical properties of α -L-iduronidase are well characterized.

Mutagenesis

Studies to assess the mutagenic potential of rhIDU have not been conducted, and there are no known interactions with DNA. Mutagenic potential would not be anticipated with Aldurazyme[®] based on the structure of the drug substance (a recombinant human glycoprotein), its impurity profile and the excipients in the final product (polysorbate 80, sodium phosphate, and sodium chloride).

Impairment of Fertility

No effects on mating and fertility parameters were observed in an intravenous fertility and general reproduction toxicity study of rhIDU in rats. There were no treatment-related effects on sperm parameters or findings on gross necropsy. There was no effect on litter parameters or treatment-related fetal effects. Male rats received rhIDU or vehicle from 28 days before cohabitation until sacrifice after 7 days of cohabitation. Female rats received rhIDU or vehicle from 15 days before cohabitation until the seventh day of gestation (DG 7); they were sacrificed on DG 21. In contrast to the once-per-week dosing in efficacy studies, the rats in this study received once daily IV doses of 0, 0.036, 0.36 or 3.6 mg/kg/day. Symptoms of anaphylaxis were observed at 0.36 mg/kg and 3.6 mg/kg and consequently all rhIDU-treated rats were treated thereafter with decreasing (with time) doses of diphenhydramine. There were no other treatment-related clinical signs, mortality or dose/treatment-related effects on male body weights or feed consumption. A significant decrease in body weight gain was observed in females at 3.6 mg/kg

on DG 0 to 8. The NOAEL determined in this study was >3.6 mg/kg males and >0.36 mg/kg in females. The maternal NOAEL determined in this study was > 0.36 mg/kg. The developmental NOAEL for rhIDU was determined to be > 3.6 mg/kg.

Single Dose Toxicity

Acute intravenous toxicity studies of rhIDU were conducted in rats and canines. Overall, there were no treatment-related findings in the acute rat and canine toxicity studies at doses up to approximately 1.7- and 10-fold, respectively, the recommended human dose of 100 U/kg (0.58/kg).

Repeated Dose Toxicity

A repeat dose intravenous toxicity study was conducted in 32 cynomolgus monkeys. rhIDU caused no significant toxicity (ophthalmic, electrocardiographic, organ weight, or macroscopic and microscopic pathology) when administered IV weekly for 26 weeks at doses up to 16.6 mg/kg. This dose is equivalent on a surface area basis to a dose of approximately 5.5 mg/kg in humans, which is approximately 10-fold the dose administered in the Phase 1/2 and Phase 3 studies (0.58 mg/kg). All of the monkeys treated with rhIDU developed antibodies in a dose-dependent manner. Antibody levels declined in approximately half of the monkeys between Weeks 13 and 26; levels rose in the remaining animals. The change in antibody levels between Weeks 13 and 26 was not dose-dependent. Only one event consistent with a mild hypersensitivity response to administration of drug was observed in this study. Edema developed at Week 4 in one male monkey in the 16.6 mg/kg dose group.

An 8-week repeat intravenous dose study in canines was undertaken to assess the possible effects of infusate diluents on the ability of the current clinical drug product to provoke anaphylactoid reactions in normal canines. None of the three different infusion formulations of rhIDU (saline, saline plus canine serum albumin, and saline with polysorbate 80) induced an anaphylactoid reaction, although transient infusion-related reactions developed during the third dose (Week 3) that resolved with subsequent infusions. The status of the cardiovascular system during these events was found to be within normal limits. All canines developed IgG antibodies in this study and fixed complement during the infusions where infusion-related reactions occurred.

Reproductive Toxicity

In an intravenous fertility and general reproduction toxicity study of rhIDU in rats, there was no reproductive toxicity at doses up to 3.6 mg/kg.

In an intravenous developmental toxicity study of rhIDU in rats, no rhIDU-treatment related effects on litter parameters were observed after ten consecutive daily infusions of rhIDU (0.036, 0.36 or 3.6 mg/kg/day). Symptoms of anaphylaxis were observed at 0.36 mg/kg/day and

consequently all rhIDU-treated rats were treated thereafter with IV doses of diphenhydramine, No mortality or other treatment-related clinical signs were observed. Body weight gains were significantly reduced on DG 10 to 12 and body weights were slightly reduced vs. controls on DG 7 to 18 at 0.36 and 3.6 mg/kg. At these doses, there was also a significant reduction in absolute and relative food consumption on DG 15 to 18. Body weights and food consumption were unaffected at 0.036 mg/kg dose.

REFERENCES

1. Belani KG, Krivit W, Carpenter BLM, Braunlin E, Buckley JJ, Liao JC et al. Children with Mucopolysaccharidosis: Perioperative Care, Morbidity, Mortality and New Findings. *J Ped Surg* 1993; 28(3):403-410.
2. Bittner V, Weiner, DH, Yusuf, S, William, JR et al. Prediction of Mortality and Morbidity with a 6-Minute Walk Test in Patients With Left Ventricular Dysfunction. *JAMA* 1993; 270 (14):1702-1707.
3. Dangel JH. Cardiovascular Changes in Children with Mucopolysaccharide Storage Diseases and Related Disorders--Clinical and Echocardiographic Findings in 64 patients. *Eur J Pediatr* 1998; 157(7):534-538
4. Guyatt GH, Sullivan MJ, Thompson PJ, Fallen EL, Pugsley SO, Taylor DW et al. The 6-Minute Walk: a New Measure of Exercise Capacity in Patients with Chronic Heart Failure. *Can Med Assoc J* 1985; 132(8):919-923.
5. Kakkis ED, McEntee MF, Schmidtchen A, Neufeld EF, Ward DA, Gompf RE et al. Long-Term and High-Dose Trials of Enzyme Replacement Therapy in the Canine Model of Mucopolysaccharidosis I. *Biochem Mol Med* 1996; 58(2): 156-167
6. Kakkis ED, Muenzer J, Tiller GE, Waber L, Belmont J, Passage M et al. Enzyme- Replacement Therapy In Mucopolysaccharidosis I. *N Engl J Med* 2001; 344(3):182-188
7. Kakkis ED, Schuchman E, He X, Wan Q, Kania S, Wiemelt S et al. Enzyme Replacement Therapy In Feline Mucopolysaccharidosis I. *Mol Genet Metab* 2001; 72(3): 199-208.
8. Leighton SE, Papsin B, Vellodi A, Dinwiddie R, Lane R. Disordered Breathing During Sleep in Patients with Mucopolysaccharidoses. *Int J Pediatr Otorhinolaryngol* 2001; 58(2):127-138
9. Neufeld E F, Muenzer J. The Mucopolysaccharidoses. In: Scriver CR, Beaudet AL, Sly WS, Valle D, editors. *The Metabolic and Molecular Bases of Inherited Disease*. New York, McGraw-Hill, 2001: 3421-3452.

10. Roubicek M, Gehler J, Spranger J. The Clinical Spectrum of α -L-Iduronidase Deficiency. *Am J Med Genet* 1985; 20(3):471-481.
11. Scott HS, Bunge S, Gal A, Clarke LA, Morris CP, Hopwood JJ. Molecular Genetics Of Mucopolysaccharidosis Type I: Diagnostic, Clinical And Biological Implications. *Hum Mutat* 1995; 6:288-302.
12. Shull RM, Kakkis ED, McEntee MF, Kania SA, Jonas AJ, Neufeld EF. Enzyme Replacement in a Canine Model of Hurler Syndrome. *Proc Natl Acad Sci USA* 1994; 91:12937-12941.
13. Wraith JE, Clarke LA, Beck M, Kolodny EH, Pastores GM, Muenzer J, Rapoport DM, Berger KI, Swiedler S, Kakkis ED, Braakman T, Chadbourne E, Walton-Bowen K, Cox GF. Enzyme replacement therapy for mucopolysaccharidosis I: A randomized, double-blind, placebo-controlled, multinational study of recombinant human α -L-Iduronidase (Aldurazyme). *J. Pediatr* 2004; 144:581-588.

PART III: CONSUMER INFORMATION

Pr **Aldurazyme**[®]
Laronidase

This leaflet is part III of a three-part "Product Monograph" published when Aldurazyme[®] was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about Aldurazyme[®]. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATIONWhat the medication is used for:

Aldurazyme[®] (laronidase) is used to treat the non-central nervous system manifestations of Mucopolysaccharidosis I (MPS I; α -L-iduronidase deficiency) in patients with a confirmed diagnosis of this disease.

What it does:

Patients with MPS I are deficient in the enzyme α -L-iduronidase. Laronidase is a form of α -L-iduronidase produced by recombinant DNA technology. Laronidase can help to treat some of the symptoms of MPS I by replacing the deficient enzyme.

When it should not be used:

Do not use Aldurazyme[®] if you are allergic to laronidase or to any ingredient of Aldurazyme[®] or component of the container.

What the medicinal ingredient is:

Laronidase

What the important nonmedicinal ingredients are:

Polysorbate 80, Sodium chloride, Sodium phosphate monobasic monohydrate, Sodium phosphate dibasic heptahydrate, Water for Injection

For a full listing of nonmedicinal ingredients see Part I of the product monograph.

What dosage forms it comes in:

Aldurazyme[®] is supplied as a sterile concentrate for solution to be used as intravenous infusion.

WARNINGS AND PRECAUTIONSSerious Warnings and Precautions

Do not use Aldurazyme[®] if you are severely allergic to any ingredients of Aldurazyme[®] or if you have experienced a severe allergic or anaphylactic reaction to laronidase.

If you are treated with Aldurazyme[®] you may experience an infusion related reaction. Infusion related reaction is defined as any related side effect occurring during the infusion or during the 3 hours following infusion. Life-threatening infusion related reactions including anaphylactic reactions have been observed in some patients during Aldurazyme[®] infusions. Reactions have included: inability to breath independently, difficulty breathing, noisy breathing, fast breathing, temporary narrowing of the

airway, partial or complete blockage of the airway, low levels of oxygen in the blood, low blood pressure, slow heartrate, and hives. Interventions have included: life saving emergency medical treatment, the use of a machine to help with breathing,, emergency access to the patient's windpipe, and hospitalization. Other treatment may include inhaled beta-adrenergic agonists to help with breathing, epinephrine as part of emergency care and intravenous corticosteroids to help fight inflammation. In clinical trials and post-marketing experience with Aldurazyme[®], approximately 1% of patients experienced severe or serious allergic reactions. In patients with MPS I, pre-existing upper airway obstruction may have contributed to the severity of some reactions. Because of the potential for severe infusion reactions, appropriate medical support should be readily available when Aldurazyme[®] is administered. Because of the potential for recurrent severe reactions, some patients may require additional observation.

Patients with an acute underlying illness (e.g. severe infections, bronchitis, wheezing/difficulty in breathing) at the time of Aldurazyme[®] infusion may be at risk for infusion-related reactions. Careful consideration should be given to your clinical status prior to administration of Aldurazyme[®].

BEFORE you use Aldurazyme[®], talk to your doctor or pharmacist if:

- If you have an acute underlying illness
- If you have had a severe allergic or anaphylactic reaction to administration of Aldurazyme[®]
- Any allergies to this drug or its ingredients or components of the container
- If you are pregnant or plan to become pregnant or are breast-feeding

INTERACTIONS WITH THIS MEDICATIONDrugs that may interact with Aldurazyme[®] include:

No formal drug/drug interaction studies have been conducted. Please inform your doctor if you use medicinal products containing chloroquine or procaine due to the potential risk of interference with the intracellular uptake of laronidase.

PROPER USE OF THIS MEDICATIONUsual dose:

The recommended dosage regimen of Aldurazyme[®] is 0.58 mg/kg body weight administered once weekly as an intravenous infusion.

Aldurazyme[®] treatment should be supervised by a physician experienced in the management of patients with MPS I or other inherited metabolic diseases.

Administration of Aldurazyme[®] should be carried out in an appropriate clinical setting where resuscitation equipment to manage medical emergencies would be readily available.

Overdose:

There is no experience with overdoses of Aldurazyme®.

Missed Dose:

If you have missed an Aldurazyme® infusion, please contact your doctor.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Like all medicines, Aldurazyme® can have side effects.

Side effects were mainly seen while patients were being given the medicine or shortly after (infusion-related reactions). The number of these reactions decreased the longer the patients were on Aldurazyme®. The majority of these reactions were of mild intensity.

In clinical trials, the most common reactions in patients 6 years and older were flushing and headache. The most common infusion associated reactions in patients less than 5 years of age were fever and chills. In an early clinical study, three patients had episodes of swelling of their mouth and breathing passage (see table below).

In post-marketing experience with Aldurazyme®, severe and serious infusion-related reactions have been reported, some of which were life threatening. The most frequently reported side effects included chills, vomiting, nausea, joint pain, diarrhea, fast heart rate, abdominal pain, blood pressure increased, and oxygen saturation decreased. Additional adverse reactions identified in the post marketing setting also include bluish color of the skin (due to lower levels of oxygen in blood), feeling cold and redness of skin. There have been a small number of reports of leakage of IV drug from the injection site into the surrounding area under the skin. However, there have been no reports that this leakage has caused severe damage to this area under the skin near the injection site.

If you exhibit such a reaction following the administration of Aldurazyme®, you should immediately contact your doctor. You may be given additional medication such as antihistamines and paracetamol to help prevent allergic-type reactions.

SERIOUS SIDE EFFECTS and (INFUSION REACTIONS), HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM		
Symptom / effect		Talk with your doctor or pharmacist
Common (occurred in <10% of patients)	Back pain, headache, joint pain, rash, abdominal pain, severe allergic reaction with airway obstruction, swelling of the mouth and breathing passage	Yes
20 Patients younger than 5 years treated for up to 12 months		
Very Common (occurred in ≥ 10% of patients)	Fever, chills, fast heart rate, increased blood pressure, decreased oxygen in the blood	Yes
Common (occurred in <10% of patients)	Heart rate increased, respiratory distress, wheezing, itching, rash	Yes

This is not a complete list of side effects. For any unexpected effects while taking Aldurazyme®, contact your doctor or pharmacist.

HOW TO STORE IT

Keep out of reach and sight of children. Store under refrigeration at 2°C to 8°C (36°F to 46°F). Do not use after the expiration date on the vial. This product contains no preservatives.

MPS I Registry:

A registry for MPS I patients has been established in order to better understand the variability and progression of MPS I disease, and to continue to monitor and evaluate treatments. You are encouraged to participate. Your participation, or your child's participation, may involve long-term follow-up. Information regarding the registry program may be found at www.MPSIregistry.com or by calling (800) 745-4447. If you are interested in participating, please contact your doctor. You can only participate in the Registry through your doctor.

SERIOUS SIDE EFFECTS and (INFUSION REACTIONS), HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM		
Symptom / effect		Talk with your doctor or pharmacist
45 Patients 6 years and older treated for up to 12 months		
Very common (occurred in ≥ 10% of patients)	Flushing, joint disease	Yes

REPORTING SUSPECTED SIDE EFFECTS

To monitor drug safety, Health Canada through the Canada Vigilance Program collects information on serious and unexpected side effects of drugs. If you suspect you have had a serious or unexpected reaction to this drug you may notify Canada Vigilance:

Online: www.healthcanada.gc.ca/medeffect

Toll-free phone: 1-866-234-2345

Toll-free fax: 1-866-678-6789

Postage Paid Mail:

Canada Vigilance Program

Health Canada

AL 0701C

Ottawa ON K1A 0K9

NOTE: Should you require information related to the management of the side effect, please contact your health care provider. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found at: www.genzyme.ca or by contacting the sponsor, Genzyme Canada Inc., at: 1-877-220-8918

This leaflet was prepared by Genzyme Canada Inc.
Date of Approval: February 12, 2010

Aldurazyme[®] is a registered trademark of BioMarin/Genzyme LLC.